Clinical Innovation -Butterfly Appliance.

Dr.Dharmesh HS, Dr.Bharathi VS

Date of Submission: 09-06-2023

Date of Acceptance: 19-06-2023

ABSTRACT: Skeletal anterior open bite can be treated with mini-screws sincethey provide an absolute anchorage to correct it through maxillarymolar intrusion. With an adequate control a bite block was used n this case to help correct the malocclusion. Thetreatment option selected for the patient depends on the esthetic and functional considerations. Treatment of anterior open bite by extrusion of the maxillary anterior teeth in a patient with good incisor show can lead to increased gingival display.TAD-based intrusion of the posterior teeth in open bite cases has been found to be considerably stable. With the introduction of TADs, orthognathic surgery could be avoided for select anterior open bite cases. This appliance made of cast metal alloy with palatal TAD's successfully used to treat the skeletal anterior open

Key words: TAD , Skeletal open bite, molar intrusion

I. INTRODUCTION:

Anterior open bite is considered to be one of themost difficult problems to treat in orthodontics. In conventionalorthodontic treatment, various treatment modalities for the correction of an anterior open bite havebeen proposed such as extrusion of the anterior teethusing intermaxillary elastics. ^{1,2}Skeletal anterior open bite is a complicated malocclusion characterized mainly by overgrowth of the maxillary and mandibular posterior dentoal veolarheights, resulting in a longer vertical facial dimension and a steeper mandibular plane. ^{4,5}With the absolute intrusion of the posterior teeth, it is possible to autorotate the mandible in a

closingcounterclockwise direction, close the open bite, andreduce the anterior facial height without the need for surgical intervention. Miniscrewshave many advantages over other various temporary anchorage devices. Miniscrews are relatively simple and easy to insert, less traumatic, stable for the optimal force, and make it possible to apply a force immediately after insertion.³The treatment mechanics of anterior open bite with posterior intrusion by using microscrew implants were effective but still require a proper retention protocol. The intrusion of the maxillary molars with miniscrews is an interesting option in selected cases of skeletal anterior open bite. The retention protocol however, should be specific in these cases.7

.....

Appliance Design:The Butterfly appliance consists of two parts a) Cobalt chromium metal casting ⁸ on the occlusal surfaces of 25,26,27 and 15,16,17 b) I shaped casting metal placed towards the palate for the engagement of the miniscrews.(Fig:2)

Cobalt chromium metal casting engaging the occlusal surfaces of 25,26,27 and 15,16,17 is given along with a separate 'I' shaped cobalt chromium metal casting for the engagement of the miniscrews in the palatal region .The cobalt chromium occlusal casting consists of two circular holes in the connection bars for the engagement of the e-chain which inturn connects to the I shaped metal flange. The I shaped metal flange consists of hooks on the open ends for connecting the Echain which inturn applies a intrusive force on the molars.



Fig 1: Intraoral patient models showing anerior open bite.



Fig 2: Butterfly appliance fabricated on the master cast.



Case Description: A male patient aged about 22 years reported to the department with the chief complaint of forwardly placed upper and lower incisors, anterior open bite, spacing i.r.t 11-12, 21-22, 22-23, rotation.i.r.t 12,14,15,24, 25,42, 33,34, 35, 44 and 45 with positive lip step, hypotonic upper lip and hyperactive mentalis muscle. Post a detailed analysis of the case it was decided to treat

the case with non-extraction line of treatment, implant assisted intrusion of the posterior teeth for correction of anterior open bite using the 'Butterfly appliance', (Fig:3& Fig:4) followed by leveling and aligning, retraction and space closure, finishing and detailing with fixed retainers given in the upper and lower arch.



Fig 3: Butterfly appliance cemented introrally and power chain used for intrusion engaged to the 'I' bar.



Fig 4: Intraoral pictures with the butterfly appliance cemented on the second premolar and first and second molars.

Advantages:

- 1. The appliance is customized according to the patients dental arch and is rigid and fixed.
- 2. Provides absolute intrusion in skeletal open bite cases.
- 3.Can be easily activated using E-chains.
- 4. Faster treatment results.

Drawback:

- 1.Posteriors cannot be bonded.
- 2. Chances of developing posterior crossbite
- 3.Cannot be used to bring about any other tooth movement except intrusion.
- 4.Cannot be used in periodontally compromised cases and patients with multiple missing posterior teeth
- 4. Needs to be prefabricated in the laboratory.

II. DISCUSSION:

Miniscrew anchorage has the advantages of being a simpler procedure, being minimally invasive, and requiring least patient cooperation. In this case we have used miniscrews along with the cast metal butterfly appliance with the I shaped metal flange to bring about true intrusion of the molar as the patient reported with skeletal openbite followed by positive lip step and hypotonic upper lip. It is noted that post intrusion of the molars the anterior open bite was reduced and a positive overjet was established with lip competency being achieved.

REFERENCES:

- [1]. Vertical elastics for correction of anterior open bite, Rinchuse DJ.J Clin Orthod. 1994;28:284.
- [2]. Cephalometric evaluation of open bite treatment with NiTiarch wires and anterior elastics. Am J Orthod DentofacialOrthop. 1999;116:555–562. Kucukkeles N, Acar A, Demirkaya AA, Evrenol B, Enacar A
- [3]. Open Bite Correction by Intrusion of Posterior Teeth with Miniscrew, Angle

DOI: 10.35629/5252-0503587589 | Impact Factorvalue 6.18| ISO 9001: 2008 Certified Journal Page 588





Volume 5, Issue 3, May - June 2023 pp 587-589 www.ijdmsrjournal.com ISSN: 2582-6018

- Orthodontist, Vol 78, No 4, 2008, Young-Chel Parka; Han-Ah Leeb; Nak-Chun Choib; Doo-Hyung Kim.
- Early treatment of hyperdivergent open-[4]. bite malocclusions. Semin Orthod. 2002;8:130-140.Buschang PH, Sankey W, English JD
- [5]. Anterioropen-bite malocclusion: longitudinal 10-years postretentionevaluation of orthodontically treated patients. AmJ Orthod. 1985;87:175-186.Lopez-Gavito Wallen TR, Little RM, Joondeph DR
- [6]. Nonextraction treatment of an open bite with microscrew implant anchorage, AJODO, 2006 Sep;130(3):391-402.Hyo-Park, Oh-Won Kwon, Jae-Hyun Sang Sung.
- Nonsurgical treatment and stability of an [7]. adult with a severe anterior open-bite malocclusionAldo Cambiano, Guilherme Janson, Diego Coelho Lorenzoni, Daniela Gamba Garib, Dino Torres Dávalos
- [8]. D-Arch: An Interdisciplinary Approach to Absolute Anchorage.Journal of Indian orthodontic society. Vol 53, Issue 1 April 15, 2019, H S Dharmesh, S V Bharathi, H Kiran.